




Contents lists available at ScienceDirect

Complementary Therapies in Clinical Practice

journal homepage: www.elsevier.com/locate/ctcp

Evaluating the efficacy of lavender aromatherapy on peripheral venous cannulation pain and anxiety: A prospective, randomized study



Tugba Karaman^{*}, Serkan Karaman, Serkan Dogru, Hakan Tapar, Aynur Sahin, Mustafa Suren, Semih Arici, Ziya Kaya

Department of Anesthesiology and Reanimation, School of Medicine, Gaziosmanpasa University, 60100, Tokat, Turkey

ARTICLE INFO

Article history:
Received 23 March 2016
Accepted 23 March 2016

Keywords:
Aromatherapy
Lavandula
Cannulation
Pain
Anxiety

ABSTRACT

Objective: This study was designed to evaluate the effectiveness of lavender aromatherapy on pain, anxiety, and level of satisfaction associated with the peripheral venous cannulation (PVC) in patients undergoing surgery.

Method: One hundred and six patients undergoing surgery were randomized to receive aromatherapy with lavender essential oil (the lavender group) or a placebo (the control group) during PVC. The patients' pain, anxiety, and satisfaction scores were measured.

Results: There was no statistically significant difference between the groups in terms of demographic data. After cannulation, the pain and anxiety scores (anxiety 2) of the patients in the lavender group were significantly lower than the control group (for $p = 0.01$ for pain scores; $p < 0.001$ for anxiety 2 scores). In addition, patient satisfaction was significantly higher in the lavender group than in the control group ($p = 0.003$).

Conclusion: Lavender aromatherapy had beneficial effects on PVC pain, anxiety, and satisfaction level of patients undergoing surgery.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

Peripheral intravenous cannulation is a mandatory phase in patients undergoing general or regional anesthesia during surgery. Although it is one of the most common invasive procedures performed by an anesthesiologist, it is often disregarded that it can lead to severe pain, anxiety, and discomfort [1,2]. However, these unfavorable feelings may cause needle phobia, which could result in the avoidance of receiving medical care [3].

Pain is described as an unpleasant sensory and emotional experience associated with actual or potential tissue damage. The nature of pain is not obvious; it is known that pain consists of physical and psychological factors [4]. Moreover, pain and emotion have been shown to have common linguistic and neurologic pathways [5,6]. Thus, not surprisingly, several studies have

presented information on the linear relationship between anxiety and acute pain [7–9].

In alternative medicine, aromatherapy is a way to use essential oils, and it is becoming increasingly popular. Research has confirmed the anxiolytic effect of aromatherapy with lavender [10,11]. However, its analgesic effect is unknown. Some studies that have evaluated pain that arises from surgery failed to identify the positive analgesic effects of lavender [12,13]. A literature search revealed that only one study [14] has evaluated the effect of aromatherapy with lavender oil on pain and anxiety during needle insertion in healthy volunteers. Nevertheless, there is no data about the effect of aromatherapy with lavender oil on pain in patients under high stress conditions, such as surgery. Thus, this study was designed to test the hypothesis that aromatherapy with lavender oil can decrease pain and anxiety and increase the satisfaction scores of patients undergoing surgery during the placement of a peripheral venous access catheter.

^{*} Corresponding author. Department of Anesthesiology and Reanimation, School of Medicine, Gaziosmanpasa University, Ali Şevki EREK Campus, School of Medicine, 60100, Tokat, Turkey.

E-mail address: drtugbaguler@hotmail.com (T. Karaman).

2. Materials and method

2.1. Patients and study design

This prospective, randomized, single-blind, parallel-group, placebo-controlled study was performed during peripheral venous cannulation in patients that underwent surgery from November 2015 to March 2016. Study approval was obtained from the Gaziosmanpasa University Clinical Research Ethics Board (15-KAEK-132, 25/08/2015). After registering at www.clinicaltrials.gov (NCT02592044), and after obtaining written informed patient consent, patients with American Society of Anesthesiologist (ASA) physical status I and II, between the ages of 18 and 65, were scheduled for elective surgery at the Gaziosmanpasa University School of Medicine Hospital. Patients with a history of anxiety disorders, preoperative pain, asthma, chronic obstructive pulmonary disease, a poor sense of smell, and allergies to lavender essential oil and anxiolytic or analgesic drugs, or who were pregnant or breastfeeding, were excluded from the study. A researcher that was not involved in the study evaluated the eligibility criteria and selected the patients that participated in the study. The patients were randomized into two groups: a lavender group and a control group, on a 1:1 ratio using a computer-generated random table. The group allocation information was concealed in a sealed opaque envelope and opened by the preoperative care room (PCR) nurse on the day of the procedure. The patient and the outcome assessor were unaware of the group allocation. However, because of lavender's odor, it was impossible to blind the patients from knowing that they were being treated with the essential oil.

2.2. Procedure

No medication was administered to the patients before the procedure. In the PCR, the patients were monitored using electrocardiography and pulse-oximetry, and their blood pressure was recorded. The PCR nurse placed either two drops of 1% lavender essential oil (*Lavandula angustifolia* oil, Talya Herbal Products, Antalya, Turkey) or pure water on a 5 × 5 cm impermeable gauze pad and asked the patients to inhale for 5 min, while in a seated position. After 5 min, the patients were asked to turn their head to the left and an 18G venous cannula (Med Devices, London, United Kingdom) was inserted into a peripheral vein on the back of the right hand. The patients also continued to inhale either the lavender essential oil or the pure water during cannulation. If cannulation was not achieved at the first attempt, the patient was excluded from the study.

2.3. Outcomes and assessment

The primary outcome of this study was to determine the patients' pain scores during the peripheral venous cannula insertion process. Pain was evaluated by the researcher, who was blind to the patient allocation group, using the 10 cm visual analog scale (VAS), 2 min after cannulation in the PCR. The secondary outcomes of the study were to determine the patients' anxiety and satisfaction levels. The same researcher measured and recorded each patient's anxiety level (anxiety 2) using the 10 cm VAS, 2 min after cannulation; patient satisfaction was measured using the 5-point Likert scale (0: worst to 4: best) 15 min after cannulation. The demographic data, age, sex, body mass index (BMI), ASA physical status, and anxiety scores (anxiety 1) of the patients at admittance to the PCR were also noted.

2.4. Sample size

In our previous study [15], the mean and standard deviation (SD) pain scores of patients during peripheral venous cannulation were found to be 3.16 ± 2.21 . We then hypothesized that aromatherapy with lavender essential oil could reduce the VAS scores by 40%. Assuming a two-sided type 1 error 0.05 ($\alpha = 0.05$) and a power of 0.80 ($\beta = 0.02$), we calculated that a minimum of 48 patients per group would be required. We increased the sample size by 10%–106 patients (53 patients per group) to account for the possibility of study drop outs.

2.5. Statistical analysis

All statistical analyses were performed using Statistical Package for the Social Sciences (SPSS) software version 20.0 (SPSS Inc., Chicago, IL, USA). Descriptive data were presented as mean (\pm SD) for the continuous variables, median (range) for the ordinal variables, or as a number (frequencies) for the categorical variables. The one-sample Kolmogorov-Smirnov test was used to detect the normality of distribution. The normally-distributed continuous variables were evaluated using independent samples t-tests. The continuous variables that were not normally distributed were compared using Mann-Whitney U tests. The categorical variables were analyzed using Pearson's chi-square or Fischer's exact tests. Analysis of the correlation among the variables was conducted using Spearman's correlation coefficients. The Wilcoxon's test was used to evaluate the pretreatment and post-treatment data; p values < 0.05 were considered to be statistically significant.

3. Results

One hundred and fifty patients that were scheduled for elective surgery were assessed for eligibility to participate in the study, and 44 of them were excluded because they did not meet the inclusion criteria, or they declined to participate. A total 106 patients were enrolled in the study; 53 patients were randomized into the lavender group and 53 patients were randomized into the control group. After being enrolled in the study, two patients in the lavender group and three patients in the control group were excluded because of protocol violations. The details for the study are shown in the consort flow diagram (Fig. 1).

The two groups were comparable in terms of the patients' demographic characteristics (age, sex, BMI)(Table 1).

The patients' mean (range) pain score was 1.94 (1–4) in the lavender group and 2.48 (1–5) in the control group. The patients' pain scores in the lavender group were significantly lower during the venipuncture process in comparison to the control group ($p = 0.01$; Fig. 2).

The anxiety scores of the patients were similar in both groups at admittance to the PCR ($p = 0.80$, Table 2). However, after aromatherapy with lavender essential oil, the anxiety scores in the lavender group were significantly lower than the control group (pure water placebo; $p < 0.001$, Table 2, Fig. 3). The pain scores showed a moderate correlation with the anxiety 2 scores when all patients evaluated (Spearman's rho, $r = 0.258$, $p = 0.009$). However, no correlation was found between the pain scores and the change (Δ anxiety) in the anxiety scores (anxiety 1 – anxiety 2) ($p = 0.233$).

The mean (range) satisfaction level of the patients was 2.29 (1–4) in the lavender group and 1.82 (0–3) in the control group, and a significant difference was observed between the groups ($p = 0.003$, Table 2). The best degree of satisfaction (VAS: 4) was found in only one patient in the lavender group, and only one patient in the control group was found to have the worst degree of satisfaction (VAS: 0). A moderate negative correlation was found

CONSORT Flow Diagram




Fig. 1. Study flow.

Table 1 Patients' characteristics.

	Lavender group	Control group	p value
Age (years)	41.51 ± 12.39	45.24 ± 10.81	0.11
Sex (F/M)	23/28	24/26	0.77
Body Mass Index (kg/m ²)	25.25 ± 2.61	26.12 ± 1.92	0.06
ASA (I/II)	36/15	34/16	0.78

Data were expressed as mean ± SD or numbers. Female/Male (F/M).

between patient satisfaction and pain and anxiety scores regarding data of the all patients. The correlation coefficients are presented in Table 3.

4. Discussion

This study demonstrated that aromatherapy with lavender essential oil has a positive effect on the pain and anxiety levels of patients undergoing elective surgery during peripheral venous cannulation. Moreover, the satisfaction of patients that received aromatherapy with lavender essential oil was significantly improved in the PCR.

Inhalation of lavender essential oil increases blood flow and decreases systolic blood pressure and galvanic skin conduction [16]. Diego et al. [17] investigated the effect of lavender and rosemary oils on electroencephalogram (EEG) activity, alertness, and mood in adults. They showed that the lavender group had increased EEG




Fig. 2. Comparison of the Pain Scores Between Groups. Data are presented as mean ± SD. *: Significantly difference, Mann-Whitney U, p = 0.01.

Table 2
The comparison of the anxiety and satisfaction scores between groups.

	Lavender group	Control group	p value
Anxiety Scores 1	2.92 ± 1.11	3.04 ± 1.07	0.80
Anxiety Scores 2	2.04 ± 1.09	2.90 ± 0.97	^a <0.001
Satisfaction Scores	2.29 ± 0.70	1.82 ± 0.80	^a 0.003

Data were expressed as mean ± standard deviation.

^a Significantly difference, Mann-Whitney U test.




Fig. 3. Changes in Anxiety Scores. Data are presented as mean ± SD. Anxiety 1 measured at admittance into the PCR, Anxiety 2 measured 2 min after cannulation*: Significantly difference, Wilcoxon test, $p = 0.053$ in control group, $p = 0.00$ in lavender group.

Table 3
The correlation coefficients between pain, anxiety 2, and satisfaction scores.

		Satisfaction score
Pain Score	Correlation Coefficient	-0.604
	Significance	^a <0.001
	N	101
Anxiety 2 Score	Correlation Coefficient	-0.328
	Significance	^a 0.001
	N	101

^a Significantly correlation, Spearman's Rho.

beta power and drowsiness, and they were more relaxed and less depressed than the control group [17]. The anxiolytic effect of lavender was also evaluated in a study that used Bispectral index scores (BIS) and the stress levels of adults; it demonstrated that aromatherapy with lavender oil significantly reduced the BIS and stress levels [14]. Preoperative anxiety may cause increased use of anesthetic drugs, longer recovery durations, and more perioperative complications, such as sleep disorders or postoperative painful recovery [18,19]. In general, benzodiazepines or narcotic analgesics are preferred medications for decreasing preoperative anxiety. Nevertheless, these drugs can lead to increased recovery durations and postoperative cognitive dysfunctions [20]. The result of our study is consistent with the literature; we showed that patient anxiety levels decreased when lavender oil aromatherapy was used in the PCR in comparison to the placebo (pure water). Complementary therapies without pharmacological drugs may be viable alternatives for the treatment of the preoperative anxiety.

While there is clear evidence about the anxiolytic effect of lavender, firm proof of its antinociceptive effect is lacking. Gedney et al. [21], designed a randomized crossover study to examine the effects of the olfactory absorption of lavender and rosemary essential oils on sensory and affective responses to experimentally-induced pain. They found that these essential oils did not produce a significant

analgesic effect [21]. In another study, the analgesic efficacy of postoperative lavender oil aromatherapy in 50 patients undergoing breast biopsy surgery was compared with patients that only received supplemental oxygen without lavender oil; it found that postoperative narcotic requirement, pain scores, and time to discharge from the postoperative care room were not significantly different between the study groups [12].

On the other hand, some studies have verified the analgesic effect of lavender oil [14,22]. One randomized controlled clinical trial documented that the pain of patients undergoing hemodialysis following needle insertion into an arteriovenous fistula decreased using lavender oil aromatherapy [22]. Kim et al. [14] showed that the pain intensity of needle insertion in healthy volunteers was minimized after aromatherapy with lavender oil. In our study, the patients' pain was alleviated by lavender oil aromatherapy. This reduction in the level of pain could have been multifactorial. Venipuncture pain can arise from different factors, including thermoreceptors, mechanoreceptors, and nociceptors, as well as somatoform disorders [23]. The aggravating effect of anxiety on acute pain is a common clinical observation. Polaghaus et al. [24] suggested that the pain modulation effect of anxiety is associated with activation changes in the entorhinal cortex of the hippocampus. Aromatherapy was also found to have an effect on hippocampal activation [25]. Our study found a moderate correlation between the pain and anxiety 2 scores, but no correlation between the pain scores and the change in the anxiety (Δ anxiety) scores. Evidently, aromatherapy may modulate the pain either directly or via anxiety.

Patient satisfaction has recently become a core concern of health care professionals, and it has been identified as being one of the most important patient outcomes. Because patient satisfaction is multidimensional, it is difficult and complex to measure. However, some helpful tools can be used to evaluate it. Five point Likert scale as a simple instrument that can be used to quantify satisfaction [26]. We used 5 point Likert scale to evaluate the patients' pain management satisfaction, and the patients in the lavender group expressed a higher level of satisfaction than the patients in the control group. The similar finding with this study is confirmed by Kim and et al. [12,14]. They presented that the lavender aromatherapy increased the patient satisfaction without pain score difference in patients after breast biopsy surgery [12]. However, in our study, there was a significantly negative correlation between the pain, anxiety, and satisfaction scores when all of the cases were assessed. The decreased pain and anxiety scores in the lavender group could play a pivotal role in this improvement.

This study has several limitations. The first is that it could not be conducted as a double-blind study due to the odor of the lavender oil. The patients in the lavender group could not be blinded to the use of that essential oil, and the patients in the control group only inhaled pure water as a placebo, which has no odor. The second, satisfaction was evaluated at the early phase using Likert scale. The

evaluation time, and also the other effecting factors like education that unsought in our study, might influence the satisfaction.

5. Conclusion

During peripheral venous access placement, lavender oil aromatherapy could be a beneficial choice for treating pain and anxiety and improving satisfaction in patients undergoing surgery.

Conflict of interest

The authors received no financial support for the study, and they declare no conflict of interest.

Acknowledgment

We would like to thank all the nursing staff who participated in this study.

References

- [1] R. Hosseinabadi, S. Biranvand, Y. Pournia, K. Anbari, The effect of acupressure on pain and anxiety caused by venipuncture, *J. Infus. Nurs.* 38 (2015) 397–405.
- [2] M. Halter, S. Lees-Mlana, H. Snooks, K.L. Koenig, K. Miller, Out-of-hospital intravenous cannulation: the perspective of patients treated by London Ambulance Service paramedics, *Acad. Emerg. Med.* 7 (2000) 127–133.
- [3] S. Wright, M. Yelland, K. Heathcote, S.K. Ng, G. Wright, Fear of needles—nature and prevalence in general practice, *Aust. Fam. Physician* 38 (2009) 172–176.
- [4] A.J. Sturgeon, A.J. Zautra, Psychological resilience, pain catastrophizing, and positive emotions: perspectives on comprehensive modeling of individual pain adaptation, *Curr. Pain Headache Rep.* 17 (2013) 317.
- [5] A. Wierzbicka, Is pain a human universal? A cross-linguistic and cross-cultural perspective on pain, *Emot. Rev.* 4 (2012) 307–317.
- [6] E. Kross, M.G. Berman, W. Mischel, E.E. Smith, T.D. Wager, Social rejection shares somatosensory representations with physical pain, *Proc. Nat. Acad. Sci. U. S. A.* 108 (2011) 6270–6275.
- [7] E.C. Carr, V. Nicky Thomas, J. Wilson-Barnet, Patient experiences of anxiety, depression and acute pain after surgery: a longitudinal perspective, *J. Int. Nurs. Stud.* 42 (2005) 521–530.
- [8] A. Tokmak, M.K. Kokanali, A.I. Güzel, U. Taşdemir, B. Akselim, N. Yılmaz, The effect of preprocedure anxiety levels on postprocedure pain scores in women undergoing hysterosalpingography, *J. Chin. Med. Assoc.* 78 (2015) 481–485.
- [9] Z.N. Kain, F. Sevarino, G.M. Alexander, S. Pincus, L.C. Mayes, Preoperative anxiety and postoperative pain in women undergoing hysterectomy. A repeated-measures design, *J. Psychosom. Res.* 49 (2000) 417–422.
- [10] S. Fayazi, M. Babashahi, M. Rezaei, The effect of inhalation aromatherapy on anxiety level of the patients in preoperative period, *Iran. J. Nurs. Midwifery Res.* 16 (2011) 278–283.
- [11] M.Y. Cho, E.S. Min, M.H. Hur, M.S. Lee, Effects of aromatherapy on the anxiety, vital signs, and sleep quality of percutaneous coronary intervention patients in intensive care units, *Evid. Based Complement. Altern. Med.* (2013) 381381.
- [12] J.T. Kim, M. Wajda, G. Cuff, D. Serota, M. Schlame, D.M. Axelrod, et al., Evaluation of aromatherapy in treating postoperative pain: pilot study, *Pain Pract.* 6 (2006) 273–277.
- [13] R. Soltani, S. Soheilipour, V. Hajhashemi, G. Asghari, M. Bagheri, M. Molavi, Evaluation of the effect of aromatherapy with lavender essential oil on post-tonsillectomy pain in pediatric patients: a randomized controlled trial, *Int. J. Pediatr. Otorhinolaryngol.* 77 (2013) 1579–1581.
- [14] S. Kim, H.J. Kim, J.S. Yeo, S.J. Hong, J.M. Lee, Y. Jeon, The effect of lavender oil on stress, bispectral index values, and needle insertion pain in volunteers, *J. Altern. Complement. Med.* 17 (2011) 823–826.
- [15] M. Suren, Z. Kaya, M. Gokbakan, I. Okan, S. Arici, S. Karaman, et al., The role of pain catastrophizing score in the prediction of venipuncture pain severity, *Pain Pract.* 14 (2014) 245–251.
- [16] Y. Saeki, M. Shiohara, Physiological effects of inhaling fragrances, *Int. J. Aromather.* 11 (2001) 118–125.
- [17] M.A. Diego, N.A. Jones, T. Field, M. Hernandez-Reif, S. Schanberg, C. Kuhn, et al., Aromatherapy positively affects mood, EEG patterns of alertness and math computations, *Int. J. Neurosci.* 96 (1998) 217–224.
- [18] T.M. Osborn, N.A. Sandler, The effects of preoperative anxiety on intravenous sedation, *Anesth. Prog.* 51 (2004) 46–51.
- [19] Z.N. Kain, L.C. Mayes, A.A. Caldwell-Andrews, D.E. Karas, B.C. McClain, Pre-operative anxiety, postoperative pain, and behavioral recovery in young children undergoing surgery, *Pediatrics* 118 (2006) 651–658.
- [20] K. Millar, A.J. Asbury, A.W. Bowman, M.T. Hosey, K. Martin, T. Musiello, et al., A randomised placebo-controlled trial of the effects of midazolam premedication on children's postoperative cognition, *Anaesthesia* 62 (2007) 923–930.
- [21] J.J. Gedney, T.L. Glover, R.B. Fillingim, Sensory and affective pain discrimination after inhalation of essential oils, *Psychosom. Med.* 66 (2004) 599–606.
- [22] M. Bagheri-Nesami, F. Espahbodi, A. Nikkhah, S.A. Shorofi, J.Y. Charati, The effects of lavender aromatherapy on pain following needle insertion into a fistula in hemodialysis patients, *Complement. Ther. Clin. Pract.* 20 (2014) 1–4.
- [23] S.J. Fetzer, Reducing venipuncture and intravenous insertion pain with eutectic mixture of local anesthetic: a meta-analysis, *Nurs. Res.* 51 (2002) 119–124.
- [24] A. Ploghaus, C. Narain, C.F. Beckmann, S. Clare, S. Bantick, R. Wise, et al., Exacerbation of pain by anxiety is associated with activity in a hippocampal network, *J. Neurosci.* 21 (2001) 9896–9903.
- [25] L. Rombola, M.T. Corasaniti, D. Rotiroti, C. Tassorelli, S. Sakurada, G. Bagetta, et al., Effects of systemic administration of the essential oil of bergamot (BEO) on gross behaviour and EEG power spectra recorded from the rat hippocampus and cerebral cortex, *Funct. Neurol.* 24 (2009) 107–112.
- [26] C.K. Ross, C.A. Steward, J.M. Sinacore, A comparative study of seven measures of patient satisfaction, *Med. Care* 33 (1995) 392–406.